
WEYL DISKS



Jānis Erdmanis, Árpád Lukács, Yuli V. Nazarov


$$\mathbf{H} = \sigma \cdot \phi + \frac{1}{2} (\phi - \phi_0) \cdot \mathbf{G} \cdot (\phi - \phi_0) + V(\mathbf{q})$$
$$[\mathbf{q}, \phi] = i, \quad \mathbf{G} = diag(g_1 > g_2 > g_3)$$



- Preferred direction for ϕ will be one with the smallest g and thus with largest separation
- Thus we expect that wave function is centered around minima of kinetic term
- For large ϕ_0 we would only see one type of spin states


QUASICLASSICAL

$$E = \pm |\phi| + \frac{1}{2}(\phi - \phi_0) \cdot \mathbf{G} \cdot (\phi - \phi_0)$$
where ϕ is a minimum for E



$\begin{array}{c|c} \textbf{QUANTUM} \\ \hline & \textbf{Easy} \\ \hline & \textbf{Orthognal} \\ \hline & \boldsymbol{\phi_0} \\ \hline & \boldsymbol{\phi_0} \\ \hline & \boldsymbol{\phi_{03}} \\ \end{array} = 1$

$$\left|\frac{\partial E_0}{\partial \phi_{01,2}}\right| = \begin{cases} V = \frac{\alpha}{|\mathbf{q}|} : \frac{8\alpha^3}{g_3} \\ V = \alpha |\mathbf{q}|^2 : e^{-\frac{1}{\sqrt{2\alpha g_3^3}}} \end{cases}$$



- Energy level splitting can be approximately calculated without Weyl term
- Ground state energy is roughly a kinetic term minima plus ground state without Weyl term
- Crossings and avoidances can be explained by symmetries

CONCLUSION

- Contact effects cause energy level splitting
- Tip of a Weyl point is unaffected in the easy direction but is smeared in orthognal ones (disc)

REFERENCES

- R.-P. Riwar, M. Houzet, J. S. Meyer & Yuli V. Nazarov, Nat. Commun. 7, 11167 (2016) and its references
- Erdmanis, J., Lukács,Á., & Nazarov, Y. V. (2018). Weyl disks: Theoretical prediction. Physical Review B, 98(24).